INTERNATIONAL JOURNAL OF PHARMACEUTICS

VOL. 93 NO. 1–3 CONTENTS 31 MAY 1993

Cited in: Current Contents (Life Sciences; Clinical Practice); International Pharmaceutical Abstracts; Biological Abstracts; Chemical Abstracts; Excerpta Medica EMBASE; PASCAL M; Science Citation Index; Current Awareness in Biological Sciences

Announcement from the Publisher v

Research Papers

Effects of vitamin E and squalene on skin irritation of a transdermal absorption enhancer, lauroylsarcosine
A. Aioi, K. Kuriyama, T. Shimizu, M. Yoshioka and S. Uenoyama (Japan) 1

The interaction potential of vigabatrin with phenytoin and carbamazepine
M.I. Al-Hassan, S.A. Bawazir, K.I. Al-Khamis, K.M. Matar and A. Tekle (Saudi Arabia) 7

Application of *p-N,N*-dimethylphenylenediamine dihydrochloride for the determination of some antiarrhythmic agents and anticoagulants
C.S.P. Sastry, T.T. Rao and A. Sailaja (India) 13

High-molecular weight polyoxyethylene as an additive in ophthalmic solutions
E. Dimitrova, S. Bogdanova, E. Minkov, N. Manolova, I. Raschkov, M. Koleva and S. Mihailova (Bulgaria) 21

Transdermal delivery of 5-fluorouracil (5-FU) through hairless mouse skin by 1-alkylaminocarbonyl-5-FU prodrugs: Physicochemical characterization of prodrugs and correlations with transdermal delivery
K.B. Sloan, J.J. Getz, H.D. Beall and R.J. Prankerd (USA) 27

The estimation of relative water solubility for prodrugs that are unstable in water
H.D. Beall, J.J. Getz and K.B. Sloan (USA) 37

Ocular delivery of the β-blocker, tilisolol, through the prodrug approach
H. Sasaki, Y. Igarashi, K. Nishida and J. Nakamura (Japan) 49

Improvement of the stability of water-in-oil-in-water multiple emulsions by the addition of surfactants in the internal aqueous phase of the emulsions
T. Ohwaki, R. Machida, H. Ozawa, Y. Kawashima, T. Hino, H. Takeuchi and T. Niwa (Japan) 61

Degradation of 4′-azidothymidine in aqueous solution
M. Brandl, R. Strickley, T. Bregante, L. Gu and T.W. Chan (USA) 75

Stabilization-oriented preformulation study of photolabile menatetrenone (vitamin K_2)
R. Teraoka and Y. Matsuda (Japan) 85

Effect of absorption promoters on the nasal absorptionof drugs with various molecular weights
A. Yamamoto, T. Morita, M. Hashida and H. Sezaki (Japan) 91

Factors affecting differences in film thickness of beads coated in fluidized bed units
R. Wesdyk, Y.M. Joshi, J. De Vincentis, A.W. Newman and N.B. Jain (USA) 101

Dissolution from disintegrating tablet: separate contributions from granules and primary drug particles considered
H.G. Ibrahim and E.-S. Sallam (Jordan) 111

Physicochemical properties of the fluoroquinolone antimicrobials. V. Effect of fluoroquinolone structure and pH on the complexation of various fluoroquinolones with magnesium and calcium ions
D.L. Ross and C.M. Riley (USA) 121

Physicochemical properties of the fluoroquinolone antimicrobials VI. Effect of metal-ion complexation on octan-1-ol-water partitioning
D.L. Ross, S.K. Elkinton, S.R. Knaub and C.M. Riley (USA) 131

An application of the *C* concept in predicting the topical efficacy of finite dose acyclovir in the treatment of cutaneous HSV-1 infections in hairless mice
P.H. Lee, M.-H. Su, A.-H. Ghanem, T. Inamori, E.R. Kern and W.I. Higuchi (USA) 139

Effects of storage on the physicochemical properties and release characteristics of progesterone-loaded poly(*l*-lactide) microspheres
Y. Aso, S. Yoshioka and T. Terao (Japan) 153

Investigations into the formation and characterization of phospholipid microemulsions. I. Pseudo-ternary phase diagrams of systems containing water-lecithin-alcohol-isopropyl myristate
R. Aboofazeli and M.J. Lawrence (UK) 161

(Contents continued on p. 260)

0378-5173(19930531)93:1/3;1-K

INTERNATIONAL JOURNAL OF PHARMACEUTICS

VOL. 93 NO. 1–3 CONTENTS 31 MAY 1993

Cited in: Current Contents (Life Sciences; Clinical Practice); International Pharmaceutical Abstracts; Biological Abstracts; Chemical Abstracts; Excerpta Medica EMBASE; PASCAL M; Science Citation Index; Current Awareness in Biological Sciences

Announcement from the Publisher v

Research Papers

Effects of vitamin E and squalene on skin irritation of a transdermal absorption enhancer, lauroylsarcosine
A. Aioi, K. Kuriyama, T. Shimizu, M. Yoshioka and S. Uenoyama (Japan) 1

The interaction potential of vigabatrin with phenytoin and carbamazepine
M.I. Al-Hassan, S.A. Bawazir, K.I. Al-Khamis, K.M. Matar and A. Tekle (Saudi Arabia) 7

Application of *p-N,N*-dimethylphenylenediamine dihydrochloride for the determination of some antiarrhythmic agents and anticoagulants
C.S.P. Sastry, T.T. Rao and A. Sailaja (India) 13

High-molecular weight polyoxyethylene as an additive in ophthalmic solutions
E. Dimitrova, S. Bogdanova, E. Minkov, N. Manolova, I. Raschkov, M. Koleva and S. Mihailova (Bulgaria) 21

Transdermal delivery of 5-fluorouracil (5-FU) through hairless mouse skin by 1-alkylaminocarbonyl-5-FU prodrugs: Physicochemical characterization of prodrugs and correlations with transdermal delivery
K.B. Sloan, J.J. Getz, H.D. Beall and R.J. Prankerd (USA) 27

The estimation of relative water solubility for prodrugs that are unstable in water
H.D. Beall, J.J. Getz and K.B. Sloan (USA) 37

Ocular delivery of the β-blocker, tilisolol, through the prodrug approach
H. Sasaki, Y. Igarashi, K. Nishida and J. Nakamura (Japan) 49

Improvement of the stability of water-in-oil-in-water multiple emulsions by the addition of surfactants in the internal aqueous phase of the emulsions
T. Ohwaki, R. Machida, H. Ozawa, Y. Kawashima, T. Hino, H. Takeuchi and T. Niwa (Japan) 61

Degradation of 4′-azidothymidine in aqueous solution
M. Brandl, R. Strickley, T. Bregante, L. Gu and T.W. Chan (USA) 75

Stabilization-oriented preformulation study of photolabile menatetrenone (vitamin K_2)
R. Teraoka and Y. Matsuda (Japan) 85

Effect of absorption promoters on the nasal absorptionof drugs with various molecular weights
A. Yamamoto, T. Morita, M. Hashida and H. Sezaki (Japan) 91

Factors affecting differences in film thickness of beads coated in fluidized bed units
R. Wesdyk, Y.M. Joshi, J. De Vincentis, A.W. Newman and N.B. Jain (USA) 101

Dissolution from disintegrating tablet: separate contributions from granules and primary drug particles considered
H.G. Ibrahim and E.-S. Sallam (Jordan) 111

Physicochemical properties of the fluoroquinolone antimicrobials. V. Effect of fluoroquinolone structure and pH on the complexation of various fluoroquinolones with magnesium and calcium ions
D.L. Ross and C.M. Riley (USA) 121

Physicochemical properties of the fluoroquinolone antimicrobials VI. Effect of metal-ion complexation on octan-1-ol-water partitioning
D.L. Ross, S.K. Elkinton, S.R. Knaub and C.M. Riley (USA) 131

An application of the *C* concept in predicting the topical efficacy of finite dose acyclovir in the treatment of cutaneous HSV-1 infections in hairless mice
P.H. Lee, M.-H. Su, A.-H. Ghanem, T. Inamori, E.R. Kern and W.I. Higuchi (USA) 139

Effects of storage on the physicochemical properties and release characteristics of progesterone-loaded poly(*l*-lactide) microspheres
Y. Aso, S. Yoshioka and T. Terao (Japan) 153

Investigations into the formation and characterization of phospholipid microemulsions. I. Pseudo-ternary phase diagrams of systems containing water-lecithin-alcohol-isopropyl myristate
R. Aboofazeli and M.J. Lawrence (UK) 161

(Contents continued on p. 260)

0378-5173(19930531)93:1/3;1-K

(Contents continued from back cover)

Stabilisation of a protein prodrug formulation: a novel approach
D.M.J. Van Schie and J.G. Winchester (UK) 177
Application of the extended Hildebrand solubility parameter treatment for optimizing reversed-phase high-performance liquid chromatography determination of pharmaceuticals
A.G. Gonzalez and A.G. Asuero (Spain) 183
Investigations on the physical state of lipid nanoparticles by synchrotron radiation X-ray diffraction
K. Westesen, B. Siekmann and M.H.J. Koch (Germany) 189
Interactions between *p*-hydroxybenzoic acid esters and hydroxypropyl-β-cyclodextrin and their antimicrobial effect against *Candida albicans*
S.J. Lehner, B.W. Müller and J.K. Seydel (Germany) 201
Enhancement of dissolution rate of nifedipine using sucrose ester coprecipitates
J.D. Ntawukulilyayo, S. Bouckaert and J.P. Remon (Belgium) 209
Dissolution behaviour of drugs from binary and ternary systems
M.D. Veiga, C. Escobar and M.J. Bernad (Spain) 215
Surfactant mediated effects in pressurized metered dose inhalers formulated as suspensions. I. Drug/surfactant interactions in a model propellant system
J.G. Clarke, S.R. Wicks and S.J. Farr (UK) 221

Notes
Chitosan hydrogel as a base for semisolid drug forms
J. Knapczyk (Poland) 233
Determination of atropine sulphate and benzalkonium chloride in eye drops by HPLC
G. Santoni, A. Tonsini, P. Gratteri, P. Mura, S. Furlanetto and S. Pinzauti (Italy) 239
Simultaneous determination of enalapril maleate and hydrochlorothiazide in tablets by derivative UV spectrophotometry and high-performance liquid chromatography
G. Carlucci, E. Di Giuseppe and P. Mazzeo (Italy) 245
The compressive to tensile strength ratio of pharmaceutical compacts
J.M. Newton, G. Alderborn, C. Nystrom and P. Stanley (UK, Sweden) 249

Author index 253

Noticeboard 257

Rapid Communication
Biodegradable delivery system for single step immunization with tetanus toxoid
R. Singh Raghuvanshi, M. Singh and G.P. Talwar (India) R1